

Profile™ 2SC

Tree and Turf Growth Regulator

ACCEPTED
FOR REGISTRATION

Aug 18, 2022

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Classified for
"RESTRICTED USE"
in New York State
under 6NYCRR Part 326

Doc id: 582548

1076.41



A GROWTH REGULATOR FOR THE REDUCTION OF TERMINAL GROWTH IN TREES, HEDGES AND SHOOT GROWTH IN TURFGRASSES.

Active Ingredient

Paclobutrazol: (±)-(R*,R*)-1-[(4-chlorophenyl)methyl]-a-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....	21.8%
Other Ingredients.....	78.2%
TOTAL	100.0%

Contains 2 pounds of active ingredient per one (1) U.S. gallon.

Keep Out of Reach of Children CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted no detalle. (If you do not understand the label, find someone to explain it to you in detail).

Refer to inside of label booklet for additional precautionary information and directions for use, including first aid and storage and disposal.

Notice: Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer* and *Misuse statements* inside label booklet. If terms are unacceptable, return at once unopened.**

Specialty Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Use. Product Must be Diluted Before Use.

EPA Reg. No. 67690-22
EPA Est. No. 067690-NC-002

FPL20190826
167601

SePRO Corporation 11550 N. Meridian St., Ste. 600, Carmel, IN 46032 U.S.A.

Net contents **1 gallon (Non-refillable)**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep Out of Reach of Children

CAUTION / PRECAUCIÓN

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist.

FIRST AID

If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC at 1-800-535-5053**.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category F on an EPA chemical resistance category selection chart.

Applicators and other handlers wear:

- Long-sleeved shirt and long pants;
- Chemical-resistant gloves made of any waterproof material (such as barrier laminate, butyl rubber, nitrile rubber, or Viton® ≥14 mils); and
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all directions for use carefully before applying this product. Use only according to label directions.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated; such as plants, soil, or water is:

- Long-sleeved shirt and long pants;
- Chemical-resistant gloves made of any waterproof material (such as barrier laminate, butyl rubber, nitrile rubber, or Viton® ≥ 14 mils); and
- Shoes plus socks.

Use Information for Growth Regulation of Trees and Hedges

Profile 2SC growth regulator is a xylem mobile plant growth regulator that slows vegetative growth by inhibiting gibberellin biosynthesis. It reduces vegetative growth, thereby reducing the volume of woody growth that must be removed from trees and hedges when trimming occurs and may extend the trimming cycle. Profile 2SC is equally effective when applied as a soil-directed spray, basal soil drench or by soil injection.

Profile 2SC can be used on listed trees and hedges found in areas to include utility rights-of-way (i.e. below transmission lines, and railroad and highway rights-of-ways); urban environments (i.e. in parks and parking lot areas, storage areas, street medians and easements, commercial/municipal building or office sites, plazas, around walkways and sidewalks, above ground planters, and around schools, hospitals, and other buildings); residential areas; other non-crop areas (i.e. industrial sites, athletic fields and other recreational areas, naturalized areas and other uncultivated non-agricultural areas, fence and property lines, and airports); and in cropped areas where the tree crops (i.e. fruits and nuts) will not be harvested within one year of application.

Symptoms of Growth Regulation: Profile 2SC demonstrates little or no phloem mobility. Activity occurs following root uptake and xylem translocation throughout the tree canopy. Symptoms of growth regulation may not be visible for up to 18 months following application. Initial effects may be observed as intense greening of foliage with no resulting phytotoxicity. At the time when normal vegetative growth extension should occur, treated trees will exhibit shortened internodes which may be somewhat thickened. Smaller leaf size and increased flowering may also be observed in some species.

Use Restrictions for Applications to Trees and Hedges

- **DO NOT** treat sugar maple trees or any other trees that are or could be tapped for sugar within one year of application.
- **DO NOT** treat nut or fruit trees that will be harvested within one year of application.
- **DO NOT** treat trees which are severely stressed or rapidly declining due to moisture, temperature, low soil fertility, or exhibit mechanical or chemical injury.
- **DO NOT** reapply this product until symptoms from previous applications begin to subside, or within 3 years of the last application, whichever comes first.
- **Chemigation: DO NOT** apply this product through any type of irrigation system.
- Trees not used for food production that are not specifically listed on this label may be treated if all other label directions are followed.

Use Precautions for Applications to Trees and Hedges

- Soil type and environmental conditions can affect the degree and longevity of growth regulation following application of Profile 2SC. Follow label instructions to increase effectiveness depending on these factors.
- For soils that are hard-to-wet, the product's mobility can be enhanced by using a nonionic, organosilicone surfactant.
- If soil around tree or hedge is heavily compacted or a high water table exists, use of a registered trunk-injectable tree growth regulator such as Mastiff™ PGR may be more satisfactory.
- Spray, basal drench and soil injection application of this product may result in localized, temporary discoloration of turfgrass immediately adjacent to the treatment site.
- Avoid spray or basal drench applications on slopes or other areas where this product or treated soil may be washed away from the base of the tree or hedge by rainfall or irrigation.
- Treatment of trees bordered by shrubs and/or herbaceous ornamentals may cause these plants to be affected if their roots extend into the treatment zone.

Application Information

Application Timing

Applications can be made throughout the year, weather permitting, except when soil is frozen or saturated with water. **NOTE:** Profile 2SC is absorbed by plant roots and translocated to the growing tissues in response to evaporative water loss (transpiration). If applications are made after fall leaf drop, product uptake will not occur until development of new leaves in the spring and resumption of evaporative water loss.

Mixing Directions

Mix 317 ml (10.7 fl. oz.) of Profile 2SC with water to make one gallon of diluted mixture. One gallon of this product will make 12 gallons of diluted mixture. To improve the handling characteristics of the diluted mixture, the addition of a suspension aid is recommended. Follow all label directions and precautions on the product label of the suspension aid.

Application Methods

Profile 2SC may be applied as a spray directed at the soil beneath the trees or shrubs, as a basal drench or by soil injection. Treat only trees and hedges that are well established in their final location. Treatment of trees less than 4 inches in diameter is not recommended.

Basal Drench and spray

Prior to application, make a 2" deep furrow around the base of the tree near the point of contact between the soil and the tree trunk (refer to Figure 1). Based on tree or hedge size and species, apply the required dose (refer to Table 1). Carefully pour the diluted product mixture into the bottom of the furrow with a graduated container/jug, with a handheld hose connected to a truck-mounted tank/hydraulic sprayer, or an applicator which provides a controlled rate of flow. To avoid possible product runoff after applying, refill the furrow with untreated soil.

Soil Injection

The diluted mixture of Profile 2SC should be injected approximately 6 inches deep. Use soil injection equipment capable of delivery at 100 to 200 psi. Injection orifices should be oriented to release the diluted product horizontally at the point of injection. The required dose should be divided evenly among injection sites spaced as uniformly as possible around the trunk. The injection sites should be positioned to release diluted product mixture as close as possible to the point of contact between soil and unthickened bark beneath the soil so that the product may be readily absorbed by the tree. Injection sites should also be located next to buttress roots (refer to Figure 2). For trees less than 6 inches in diameter at breast height (DBH), use at least 4 evenly spaced injection sites per tree.




Figure 1. Placement of Profile 2SC as spray or basal drench'




Figure 2. Placement of Profile 2SC as a soil injected treatment

Application Rates

TABLE 1 Application Rate Ranges for Diluted Mixture of Profile 2SC for Treatment of Various Tree and Hedge Species[†]		
50 to 100 ml (1 to 2 grams a.i.) per inch DBH^{††}	100 to 150 ml (2 to 3 grams a.i.) per inch DBH^{††}	150 to 200 ml (3 to 4 grams a.i.) per inch DBH^{††}
Sweetgum Australian Pine	Basswood Boxelder Elm(s) Eucalyptus Hackberry Littleleaf Linden Locust Maple(s) Oak(s) Sassafrass	Ash Bradford Pear Sycamore Tuliptree Weeping Willow Ficus ^{†††}

[†] For acidic soils (low pH), soils with high organic matter content or soils with high clay content, use the higher rate, not to exceed 4 grams per inch DBH.

^{††} Diameter at breast height

^{†††} Not for use in California

TABLE 2
Amount Of Profile 2SC Diluted Mixture Required For Specific Application Rates And Tree and Hedge Sizes

DBH† (Inches)	50 ml per inch DBH†	100 ml per inch DBH†	150 ml per inch DBH†	200 ml per inch DBH†
	Total ml Required	Total ml Required	Total ml Required	Total ml Required
4	200	400	600	800
5	250	500	750	1,000
6	300	600	900	1,200
7	350	700	1,050	1,400
8	400	800	1,200	1,600
9	450	900	1,350	1,800
10	500	1,000	1,500	2,000
11	550	1,100	1,650	2,200
12	600	1,200	1,800	2,400
13	650	1,300	1,950	2,600
14	700	1,400	2,100	2,800
15	750	1,500	2,250	3,000
16	800	1,600	2,400	3,200
17	850	1,700	2,550	3,400
18	900	1,800	2,700	3,600
19	950	1,900	2,850	3,800
20	1,000	2,000	3,000	4,000
21	1,050	2,100	3,150	4,200
22	1,100	2,200	3,300	4,400
23	1,150	2,300	3,450	4,600
24	1,200	2,400	3,600	4,800
25	1,250	2,500	3,750	5,000
26	1,300	2,600	3,900	5,200
27	1,350	2,700	4,050	5,400
28	1,400	2,800	4,200	5,600
29	1,450	2,900	4,350	5,800
30	1,500	3,000	4,500	6,000
31	1,550	3,100	4,650	6,200
32	1,600	3,200	4,800	6,400
33	1,650	3,300	4,950	6,600
34	1,700	3,400	5,100	6,800
35	1,750	3,500	5,250	7,000
36	1,800	3,600	5,400	7,200

†diameter at breast height

Use Information for Growth Regulation of Turfgrass

Profile 2SC growth regulator is a plant growth regulator (PGR) which reduces leaf blade length and stem internode elongation in turfgrasses resulting in a more compact growth form for up to two (2) months after application. Growth regulation results from suppression of gibberellic acid biosynthesis. Under normal growing conditions root growth and lateral expansion of turf are not affected. An appropriate fertility program for the desired turf species should be followed in conjunction with this product's applications to provide best turfgrass enhancement and reduce potential for discoloration. Broadcast treatments should be made on medium to high quality turfgrass areas of uniform species composition. Turf containing significant amounts of coarse textured species such as tall fescue, orchardgrass, timothy, dallisgrass, etc., may respond unevenly to treatment.

Profile 2SC can be used on turfgrass grown in the following areas: golf courses (tees, fairways, greens, and rough), athletic fields, commercial and industrial areas, parks, and boulevards.

NOTICE TO USER: The rates indicated may need to be adjusted within the approved rate ranges on this label to achieve the desired level of growth regulation on turfgrass species. Turfgrass response may vary within turfgrass species due to the large number of cultivars and varieties available. The specified rate ranges permit the users to adjust the application rate to best address the growth conditions of the turfgrass being treated. Total applications must not exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year. Neither the manufacturer nor seller has determined if this product can be used safely or effectively on turfgrass species not mentioned on this label. For turfgrass species not listed on this label the user should apply this product to a small test area to determine growth response and desired level of growth regulation prior to large scale applications.

Use Restrictions for Applications to Turfgrass

- **DO NOT** use on residential turf.
- **DO NOT** apply to putting greens other than predominantly bentgrass.
- **DO NOT** apply to overseeded bermudagrass putting greens.
- **DO NOT** apply to sod farms, turfgrasses grown for seed, including plants or plant materials grown for sale or research purposes.
- **DO NOT** apply to turf within rights-of-ways.
- **DO NOT** use on areas to be cultivated for food or food crops or to be resown with grasses within two years of treatment.
- **DO NOT** apply during prolonged periods of temperature (heat or cold) or moisture stress. Also avoid applications during periods of extreme disease and insect pressure.
- **DO NOT** apply to saturated soils or when a significant moisture event is anticipated. This product may accumulate in low lying areas and cause prolonged and excessive growth regulation in those areas.
- **DO NOT** apply to areas where *Poa annua* is the desired turfgrass species or areas that contain >70% *Poa annua*.
- **DO NOT** apply to turf used for livestock production.
- **DO NOT** graze treated areas or harvest for forage or hay.
- **DO NOT** apply more than 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year.
- **Chemigation:** **DO NOT** apply through any type of irrigation system.
- Delay applications to newly seeded turfgrasses until turf is well established and actively growing.
- **DO NOT** apply until 4 weeks after turfgrass sprigging or laying sod. Turfgrass should be well established and actively growing prior to application.

Use Precautions for Applications to Turfgrass

- Additional turfgrass growth regulation may occur when product is tank mixed or used in conjunction with demethylation inhibitor (DMI) or sterol inhibiting fungicides.
- Application of an oxadiazon-based preemergence product should be spaced at least 4 weeks apart from a Profile 2SC application on putting greens.

Application Timing

Spring applications should be made after resumption of active seasonal growth of turfgrass. The final application of the season should be timed at least 4 weeks before the onset of inactive grass growth or winter dormancy. Applications to overseeded turfgrasses in dormant bermudagrass stands should be completed 4 weeks prior to expected bermudagrass green-up.

Irrigation

Profile 2SC is primarily root absorbed. After Profile 2SC, water-in within 24 hours to limit surface movement but not to the point of runoff, to limit surface movement, but not to the point of runoff. To prevent product runoff, time applications to allow for watering-in and maximum absorption into treated turf prior to a rain event.

Turf Color and Post-Treatment Turf Management:

Treated turfgrass may appear darker green in color. This color change, which appears 1 to 2 weeks after treatment, may persist an additional 3 to 8 weeks depending upon rate used and species treated. Treated areas should be managed to encourage the growth of a healthy vigorous turf.

***Poa annua* (Annual Bluegrass) Conversion to Perennial Turfgrasses:**

Applications of Profile 2SC followed by management practices designed to encourage vigorous growth of perennial turfgrasses can reduce the *Poa annua* (annual bluegrass) competition in cool season turf and increase the cover of more desirable perennial species. *Poa annua* is more sensitive to this product's treatments and is therefore more strongly suppressed than perennial turfgrass species. Discoloration of *Poa annua* can be expected from treatments that provide a desired level of growth regulator activity in perennial grass species. This effect becomes visible 7 to 10 days after treatment and lasts 3 to 6 weeks. The degree of discoloration will be proportional to the *Poa annua* composition of the turf. Product application in conjunction with soluble nitrogen fertilizers will also minimize discoloration. Application timing, rate recommendations, and precautions for perennial grass conversion through selective reduction of *Poa annua* are provided in the *Poa annua* (Annual Bluegrass) Conversion to Perennial Turfgrasses section of this label. Total applications must not exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product /acre/year.

Application Directions

Mixing Directions

Add Profile 2SC to a spray tank half filled with clean water. Allow sufficient mixing time to ensure complete dispersion and mixing. Finish filling the spray tank. Continue vigorous agitation throughout the spraying operation to ensure uniform application while avoiding overlaps and skips. Product should be applied using a boom-type sprayer with bypass and/or mechanical agitation calibrated to deliver 40 to 200 gallons/acre of spray solution (1 to 4.6 gallons/1,000 ft²). In-line strainers and nozzle screens should be 50 mesh or larger. The use of a coloring agent to mark areas already sprayed is suggested for uniform application without skips and overlaps. Performance may be improved with the addition to the spray mix of a readily available nitrogen (N) source at 0.125 to 0.5 lb N/1,000 ft² or iron (Fe) at suggested label rates.

RATE RECOMMENDATIONS FOR GROWTH REDUCTION OF PERENNIAL TURFGRASS SPECIES – MULTIPLE APPLICATION PROGRAM

Cool Season Turfgrasses

Golf courses

- Bentgrass (Golf Course Fairway Type Turf)

Multiple application program: Apply Profile 2SC to fairway height bentgrass at a rate of 16 to 32 fl. oz./acre in early spring following resumption of active growth. Repeat applications of 8 to 32 fl. oz. product/acre should be made at 2 to 6 week intervals until late summer or early fall; or repeat applications of 8 to 32 fl. oz. product/acre can be made within the same growing season as long as the turf is actively growing. Turfgrass species, growth rate, and use rate will dictate the timing of repeat applications – not to exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year.

- Bentgrass Putting Greens

Multiple application program: Apply Profile 2SC to bentgrass as part of an overall greens management program. An initial application should be made after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Apply at a rate of 4 to 16 fl. oz. product/acre. Repeat applications of 4 to 16 fl. oz. product/acre should be made at 2 to 4 week intervals. Total applications must not exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year.

Golf courses, athletic fields, commercial and industrial areas, parks, and boulevards

- Kentucky Bluegrass and Perennial Ryegrass

Multiple application program: Apply Profile 2SC to Kentucky bluegrass and perennial ryegrass at a rate of 16 to 32 fl. oz. product/acre in early spring following resumption of active growth. Repeat applications of 16 to 32 fl. oz. product/acre should be made at 4 to 6 week intervals until late summer or early fall. Total applications must not exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year. NOTE: Certain varieties of Kentucky Bluegrass are very sensitive to this product. It is recommended that this product be applied to a small test area to determine growth response and desired level of growth regulation prior to large scale applications.

Warm Season Grasses

Golf courses, athletic fields, commercial and industrial areas, parks, and boulevards

- Tifway (419), TifSport, and GN-1 Bermudagrass

Multiple application program: Apply Profile 2SC to Tifway (419), TifSport, and GN-1 Bermudagrass at a rate of 32 to 48 fl. oz. product/acre when turfgrass has completely recovered from winter dormancy and is growing vigorously. Repeat applications of 32 to 48 fl. oz. product/acre should be made at 4 to 8 week intervals until late summer or early fall. Total applications must not exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year.

TABLE 3

Rate Ranges for Growth Regulation of Perennial Turfgrass Species Using a Multiple Application Program.

Turfgrass Species	Fl. oz. Profile 2SC / A	Lbs a.i./A	Treatment Interval¹
Cool Season Turfgrasses			
Bentgrass (golf course fairway)	16 to 32	0.25 to 0.5	2 to 6 weeks
Bentgrass putting greens	4 to 16	0.063 to 0.25	2 to 4 weeks
Kentucky Bluegrass/ Perennial Ryegrass	16 to 32	0.25 to 0.5	4 to 6 weeks
Warm Season Turfgrasses			
Tifway, TifSport, and GN-1 Bermudagrass	32 to 48	0.5 to 0.75	4 to 8 weeks

¹ Repeat applications can be made within the same growing season – not to exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year. When conducting repeat applications, use shorter treatment intervals when making applications at the lower end of the rate range; use longer treatment intervals when making applications at the higher end of the rate range.

POA ANNUA (ANNUAL BLUEGRASS) CONVERSION TO PERENNIAL TURFGRASSES

Multiple Application Program

A multiple application perennial grass conversion program using this product provides *Poa annua* suppression and cool season turf conversion. This program provides a gradual perennial grass conversion reducing *Poa annua* populations over one to several growing seasons. To maximize seedling establishment, do not apply product six (6) weeks prior to and/or 2 weeks after interseeding or overseeding of perennial turfgrasses.

Golf courses

- **Bentgrass (Golf Course Fairway type turf)**

Apply Profile 2SC to fairway height bentgrass at a rate of 16 to 32 fl. oz. product/acre in early spring following resumption of active growth of the grass. Repeat applications of 8 to 32 fl. oz. product/acre should be made at 2 to 6 week intervals until late summer or early fall. Total applications must not exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year. Normal management practices such as fertilization, aeration and interseeding/overseeding will encourage growth of bentgrass.

- **Bentgrass Putting Greens**

Because annual turfgrass species such as *Poa annua* are more strongly regulated by this product, careful product use on putting greens with a high percentage of *Poa annua* can shift the competitive balance between bentgrass and *Poa annua* to favor bentgrass. Follow normal management practices such as fertilization, aeration and interseeding/overseeding that encourages growth of bentgrass. Product use on bentgrass greens may increase putting speed without reducing the height of cut.

- **Bentgrass Greens with less than 50% *Poa annua* (annual bluegrass):** Apply Profile 2SC to bentgrass as part of an overall greens management program. An initial application should be made in the spring months after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Apply at a rate of 8 fl. oz. product/acre. Repeat applications of 8 to 16 fl. oz. product/acre should be made at 2 to 4 week intervals through early fall. Total applications must not exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year.
- **Bentgrass Greens with 50 to 70% *Poa annua* (annual bluegrass):** Apply Profile 2SC to bentgrass greens at a rate of 4 to 8 fl. oz. product/acre in the spring months after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Repeat applications of 4 to 16 fl. oz. product/acre should be made at 2 to 4 week intervals through early fall. Total applications must not exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product/acre/year.

Golf courses, athletic fields, commercial and industrial areas, parks, and boulevards

- **Kentucky bluegrass, Perennial ryegrass**

Apply Profile 2SC to Kentucky bluegrass and perennial ryegrass at a rate of 16 to 32 fl. oz. product/acre in early spring following resumption of active growth of the grass. Repeat applications of 16 to 32 fl. oz. product /acre should be made at 4 to 6 week intervals until late summer or early fall. Total applications must not exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product / acre/year. Normal management practices such as fertilization, aeration and interseeding/overseeding will encourage growth of bluegrass and/or ryegrass. NOTE: Certain varieties of Kentucky Bluegrass are very sensitive to this product. It is recommended that product be applied to a small test area to determine growth response and desired level of growth regulation prior to large scale applications

TABLE 4

Rate Ranges for *Poa annua* (Annual Bluegrass) Conversion to Perennial Turfgrasses Using a Multiple Application Program

Turfgrass Species	% <i>Poa annua</i>	Initial spring application ¹		Repeat applications ¹		
		Fl. oz. Profile 2SC/A	Lbs a.i./A	Fl. oz. Profile 2SC/A	Lbs a.i./A	Treatment Interval ^{1†}
Bentgrass (golf course fairway)	0 to 70%	16 to 32	0.25 to 0.5	8 to 32	0.125 to 0.5	2 to 6 weeks
Bentgrass Putting Greens	0 to -50%	8	0.125	8 to 16	0.125 to 0.25	2 to 4 weeks
Bentgrass Putting Greens	50 to 70%	4 to 8	0.063 to 0.125	4 to 16	0.063 to 0.25	2 to 4 weeks
Kentucky Bluegrass/ Perennial Ryegrass Fairways	0 to 70%	16 to 32	0.25 to 0.5	8 to 32	0.125 to 0.5	4 to 6 weeks

¹ Apply in early spring following resumption of active growth of the grass. Fall applications should be discontinued 4 weeks before the onset of inactive grass growth or winter dormancy.

[†] Repeat applications can be made within the same growing season – not to exceed 2.0 lbs a.i./acre/year or 128 fl. oz. product /acre/year. When conducting repeat applications, use shorter treatment intervals when making applications at the lower end of the rate range; use longer treatment intervals when making applications at the higher end of the rate range.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these cannot be used according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

Container Handling:

Nonrefillable Container (rigid, 5 gallons or less): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer the container for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid, larger than 5 gal): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, treatment area, or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty Disclaimer: SePRO Corporation warrants that this product conforms to the chemical description on the product label. Testing and research have also determined that this product is reasonably fit for the uses described on the product label. To the extent consistent with applicable law, SePRO Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and any such warranties are expressly disclaimed.

Misuse: Federal law prohibits the use of this product in a manner inconsistent with its label directions. To the extent consistent with applicable law, the buyer assumes responsibility for any adverse consequences if this product is not used according to its label directions. In no case shall SePRO Corporation be liable for any losses or damages resulting from the use, handling or application of this product in a manner inconsistent with its label.

For additional important labeling information regarding SePRO Corporation's Terms and Conditions of Use, Inherent Risks of Use and Limitation of Remedies, please visit <http://seprolabels.com/terms> or scan the image below.



©Copyright 2022 SePRO Corporation

™Profile is a trademark of SePRO Corporation

™Mastiff is a trademark of ArborSystems



Profile™ 2SC

Tree and Turf Growth Regulator

A GROWTH REGULATOR FOR THE REDUCTION OF TERMINAL GROWTH IN TREES, HEDGES AND SHOOT GROWTH IN TURFGRASSES.

Active Ingredient

Pacllobutrazol: (±)-(R*,R*)-8-[[4-chlorophenyl)methyl]-α-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....21.8%

Other Ingredients.....78.2%

TOTAL.....100.0%

Contains 2 pounds of active ingredient per one (1) U.S. gallon.

Keep Out of Reach of Children CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted no detalle. (If you do not understand the label, find someone to explain it to you in detail).

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Refer to label booklet for additional precautionary information and Directions for Use, including First Aid and Storage and Disposal.

Notice: Read the entire label before using. Use only according to label directions. Before buying or using this product, read *Warranty Disclaimer and Misuse statements* inside label booklet. If terms are unacceptable, return at once unopened.

Specialty Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Use. Product Must be Diluted Before Use

EPA Reg. No. 67690-22
EPA Est. No. 067690-NC-002

FP20190826
167601

SePRO Corporation
11550 N. Meridian St., Ste. 600, Carmel, IN 46032 U.S.A.

Tree and Turf Growth Regulator

1076.41

FIRST AID

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these cannot be used according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

Container Handling:

Nonrefillable Container (rigid, 5 gallons or less): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer the container for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid, larger than 5 gal): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

See attached booklet for complete container disposal directions including triple rinse instructions.

Net Contents 1 gallon (Non-Refillable)



8

PEEL FILM HERE